MBS2006479

12th Edition (Revised in Aug, 2016)

Biotin-Linked Polyclonal Antibody to Immunoglobulin G2 (IgG2) Organism Species: Homo sapiens (Human) Instruction manual

White Courte Cou

### [PROPERTIES]

Source: Antibody labeling

Host: Rabbit

Purification: Antigen-specific Affinity Chromatography.

Label: Biotin Traits: Liquid

Concentration: 0.5mg/ml

**UOM**: 20μg(40μl)

Applications: WB; IHC; ICC.

### [ IMMUNOGEN ]

Immunogen: Native Immunoglobulin G2 (IgG2).

# [ APPLICATIONS ]

Western blotting: 0.5-2µg/mL;1:250-1000 Immunohistochemistry: 5-20µg/mL;1:25-100 Immunocytochemistry: 5-20µg/mL;1:25-100

S.COLL Optimal working dilutions must be determined by end user.

# [FORMULATION]

Form & Buffer: Supplied as solution form in 0.01M PBS, pH7.4, containing 0.05% Proclin-300, 50% glycerol. •

# [ STORAGE AND STABILITY ]

Storage: Avoid repeated freeze/thaw cycles.

Store at 4°C for frequent use.

Aliquot and store at -20°C for 12 months.

Stability Test: The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were

observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

MMM INVERSORY CE. CORP.