

PRECISION

Intra-assay Precision (Precision within an assay): CV%<8%


Three samples of known concentration were tested twenty times on one plate to assess.

Inter-assay Precision (Precision between assays): CV%<10%

Three samples of known concentration were tested in twenty assays to assess.

TYPICAL DATA

These standard curves are provided for demonstration only. A standard curve should be generated for each set of samples assayed.



pg/ml	OD1	OD2	Average	Corrected
4000	2.171	2.089	2.130	1.978
2000	1.530	1.483	1.507	1.355
1000	0.925	0.946	0.936	0.784
500	0.578	0.565	0.572	0.420
250	0.352	0.367	0.360	0.208
125	0.245	0.253	0.249	0.097
62.5	0.203	0.208	0.206	0.054
0	0.154	0.150	0.152	

LOD

15.6 pg/ml.

LINEARITY

To assess the linearity of the assay, samples were spiked with high concentrations of human HSPG2 in various matrices and diluted with the Sample Diluent to produce samples with values within the dynamic range of the assay.

	Sample	Serum(n=4)
1:1	Average %	102
	Range %	89-107
1:2	Average %	98
	Range %	91-102
1:4	Average %	94
	Range %	84-98
1:8	Average %	106
	Range %	96-110

RECOVERY

The recovery of human HSPG2 spiked to levels throughout the range of the assay in various matrices was evaluated. Samples were diluted prior to assay as directed in the Sample Preparation section.

Sample Type	Average % Recovery	Range
Serum (n=5)	103	93-106
EDTA plasma (n=4)	103	94-107